



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Solution by C. E. WHITE, Vanderbilt University, Nashville, Tenn.

$\log^n x = \log \log \log \dots (n \text{ times}) x$ .

$$\therefore d \log^n x = \frac{d(\log^{n-1} x)}{\log^n x} = \frac{d \log^{n-2} x}{\log^n x \log^{n-1} x} = \dots = \frac{dx}{\log^n x \log^{n-1} x \log^{n-2} x \dots \log x \cdot x}$$

Also solved by J. Scheffer. Some of our readers misinterpreted the meaning of the notation. It should be remembered that the notation means the log of the log of the log, etc.,  $n$  times, of  $x$ . See Byerly's *Integral Calculus*, 2d Ed., p. 2. ED. F.

251. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Find in terms of  $x$  the functions  $c_1 x$  and  $c_2 x$  defined, respectively, by the relations

$$\begin{aligned} (a) \quad x \log(c_1 x) &= c_1 x \log x, \\ (b) \quad x \log x &= c_2 x \log(c_2 x). \end{aligned}$$

Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

(a) We may write (a) thus,  $\log(c_1 x)^x = \log x^{c_1 x}$ .

$$\therefore (c_1 x)^x = x^{c_1 x}, \text{ or } c_1 x = x^{c_1} = 1 + c_1 \log x$$

$$+ \frac{c_1^2}{2!} (\log x)^2 + \dots + \frac{c_1^n}{n!} (\log x)^n + \dots$$

(b) Similarly, we may write (b),  $\log x^x = \log(c_2 x^{c_2 x})$ .

$$\therefore x^x = (c_2 x)^{c_2 x}, \text{ or } c_2 x = x^{1/c_2} = 1 + \frac{1}{c_2} \log x + \frac{1}{2! c_2^2} (\log x)^2 + \dots + \frac{1}{n! c_2^n} (\log x)^n + \dots$$

Also solved by J. Scheffer, C. E. White, and V. M. Spunar. Mr. Spuhar, in his solution, used the calculus.

## MECHANICS.

131. Proposed by F. P. MATZ.

If the distribution of weight on the foundations of a building is  $W$  lb./ (feet)<sup>2</sup>, the foundation must be sunk  $D = (W/w) \tan 4(\frac{1}{2}\pi - \frac{1}{2}\psi)$  feet deep in earth of density  $w$  lb./ (feet)<sup>2</sup> and angle of repose  $\psi$ .

Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Let  $E$  represent the weight of a portion of a horizontal stratum of earth which is displaced by the foundation of a structure,  $S$  the utmost weight of that structure consistent with the power of the earth to resist displacement,  $\psi$  the angle of repose of the earth. Then  $S/E = [(1 + \sin \psi) / (1 - \sin \psi)]^2$  (see paper "On Stability of Loose Earth," read before the Royal Society on the 19th of June, 1856, and published in the *Philosophical Transactions* for that year). In the problem,  $E = Dw$ ,  $S = W$ .

$$\therefore \frac{Dw}{W} = \left( \frac{1 - \sin \psi}{1 + \sin \psi} \right)^2 = \left( \frac{1 - \cos(\frac{1}{2}\pi - \psi)}{1 + \cos(\frac{1}{2}\pi - \psi)} \right)^2$$